




# *Carex qingyuanensis* (Cyperaceae), a new species from Guangdong, China

Ya-Li Li<sup>1,2</sup>, Shuang-Wen Deng<sup>1</sup>, Jin-Chu Luo<sup>1</sup>, Ming-Xia Li<sup>1,3</sup>, Li-Ting Zou<sup>1</sup>, Qiu-Gen Zeng<sup>4</sup>, Hong-Feng Chen<sup>1</sup>

<sup>1</sup> Key Laboratory of Plant Resources Conservation and Sustainable Utilization, Chinese Academy of Sciences, Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou 510650, China

<sup>2</sup> University of Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup> Zhongkai University of Agriculture and Engineering, Guangzhou 510225, China

<sup>4</sup> Qingyuan Forestry Bureau, Qingyuan 511515, China

Corresponding author: Hong-Feng Chen (h.f.chen@scbg.ac.cn)

## Abstract

*Carex qingyuanensis*, a new species of Cyperaceae from Guangdong Province, China, is described and illustrated. The new species is morphologically similar to *Carex peliosanthifolia* F. T. Wang & Tang ex P. C. Li, but it can be distinguished by the racemose inflorescence branches appearing single (rarely binate or ternate) (vs. binate or ternate), one (rarely two or three) (vs. 1–3) spiked, male part of linear-cylindrical spikes much longer than the female part (vs. just male part short-cylindrical and slightly longer than female part), style base thickened (vs. not thickened) and perigynium horizontally patent with a short (vs. long and excurved) beak. Phylogenetic analysis, based on the two nuclear DNA regions (ETS 1f and ITS) and three chloroplast DNA regions (*matK*, *ndhF* and *rps16*), suggests that the new species belongs to sect. *Siderostictae* s.s. of subg. *Siderosticta* and shows a closer phylogenetic relationship to *Carex scaposa* C. B. Clarke.

**Key words:** *Carex* sect. *Siderostictae* s.s., morphology, new species, taxonomy



Academic editor: Clifford Morden

Received: 22 December 2023

Accepted: 3 April 2024

Published: 25 April 2024

**Citation:** Li Y-L, Deng S-W, Luo J-C, Li M-X, Zou L-T, Zeng Q-G, Chen H-F (2024) *Carex qingyuanensis* (Cyperaceae), a new species from Guangdong, China. *PhytoKeys* 241: 155–168. <https://doi.org/10.3897/phytokeys.241.117734>

Copyright: © Ya-Li Li et al.

This is an open access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0).

## Introduction

*Carex* Linnaeus with ca. 2000 species, is one of the largest angiosperm genera and is distributed almost globally (Dai et al. 2000; Frodin 2004; Global *Carex* Group 2016). The rapid advancement of sequencing technology has significantly enhanced our comprehension of the relationships within *Carex*. According to the latest phylogenetic study, Villaverde et al. (2020) demonstrated there were six main lineages in *Carex*, identified as the *Siderostictae*, *Schoenoxiphium*, *Unispicate*, *Uncinia*, *Vignea* and *Core Carex* clades. Despite advancements, the phylogenetic relationships in *Carex* are not yet sufficient clear for a global reclassification of the genus within a Linnean infrageneric (sectional) framework. Because of this, the Global *Carex* Group (2021) employed a hybrid approach, utilizing both informally named clades and formally named sections. Their classification system represents the currently understood diversity of *Carex* lineages, encompassing six

subgenera, 62 formally designated Linnean sections, and 49 informal groups. In the recently published volume of the *Flora of China* (Dai et al. 2010), a total of 527 species were documented. Since then, nearly 40 new species and two subspecies have been described from China (Jin et al. 2004; Song et al. 2008; Jin 2009; Su 2009a, 2009b; Weng et al. 2009; Jin and Zheng. 2010; Jin et al. 2011, 2012a, 2012b; Zhao et al. 2011; Wang et al. 2012; Yu et al. 2012; Deng 2014; Yang et al. 2014; Jin and Chen 2015; Yang et al. 2015a, 2015b; Yang et al. 2016; Jin 2017; Yang et al. 2017; Lu and Jin 2018; Zhang et al. 2018; Lu et al. 2020; Yang and Liu 2020; Zhang et al. 2021; Li et al. 2022; Lu and Jin 2022; Lu et al. 2022). The new species were mainly discovered in the Provinces of Zhejiang, Guangxi and Anhui.

Many studies provide substantial evidence supporting *Carex* subg. *Siderosticta* as the sister lineage to the remaining taxa within *Carex* (Waterway et al. 2009; Starr et al. 2015; Global Carex Group 2016, 2019; Uzma Jiménez-Mejías et al. 2019; Villaverde et al. 2020). The clade exhibits considerable variability and has a narrow distribution confined primarily to East and Southeast Asia (Global Carex Group 2021).

*Carex* sect. *Siderostictae* s.s. includes 27 species, in three traditional sections: *Hemiscaposae* (12 species), *Siderostictae* s.l. (13 species) and *Surculosae* (2 species). However, these sections are not supported by the latest phylogenetic hypotheses (Global Carex Group 2021). All species in the traditional sect. *Siderostictae* s.l. and some species in the traditional sect. *Hemiscaposae* are pink-red at the base of the plant, leaves or bracts (Global Carex Group 2021). *Carex* sect. *Hemiscaposae* is unique in *Carex* for its androgynous inflorescence units, leafless pseudolateral culms and pseudopetiolate leaves. The leaves of these species are much wider (up to 12 cm) and show high similarities to *Curculigo* Gaertn (Hypoxidaceae) (Starr et al. 2015; Ford et al. 2017). Moreover, the section is noted especially for its unprecedented variety of inflorescences, ranging from simple to compound (Ford et al. 2017). Eight species have been described within sect. *Hemiscaposae*, primarily found in broadleaf evergreen forests in southern-central and southeast China (including Taiwan) (Nelmes 1955; Raymond 1959; Yang and Chen 2005; Lunkai et al. 2010). Despite the relatively small number of species, their classification is challenging due to subtle characteristic differences (Ford et al. 2017). *Carex* sect. *Siderostictae* s.s. is differentiated from the closely related sect. *Hypolytroides* by the presence of bisexual spikes and leafless fertile culms that seem to emerge laterally from leaf rosettes (Global Carex Group 2021).

During a botanical survey of Bijia Mountain Forest Farm which is located in Qingxin District, Qingyuan City, Guangdong Province, China, covering an area of 2646.11 ha, with an approximate elevation of 1000 m, a new species of *Carex* in sect. *Siderostictae* s.s. (traditionally placed in sect. *Hemiscaposae*) (subg. *Siderosticta*) was discovered in the forest on slopes. A detailed description of the species is provided below.

## Materials and methods

The material of this new species was collected during the botanical survey conducted at Bijia Mountain Forest Farm. In order to conduct a thorough morphological comparison and phylogenetic analysis, samples of similar species *Carex peliosanthifolia* and *Carex scaposa* were collected from Shangguchen, Jinxiu County, Guangxi Province, China (110°8'36.69"E, 23°53'42.15"N).

A total of 21 samples representing two sections and three clades were used for molecular phylogenetic analysis, based on the updated infrageneric classification of *Carex* (Global Carex Group 2021) and the sequencing available on GenBank. These sections and clades included Sect. *Siderostictae* s.s. (13 species), Sect. *Hypolytroides* (two species), Setigera Clade (two species), Decora Clade (two species), Esquiroliana Clade (one species), and *Eriophorum vaginatum* (Table 1). Additionally, this study provided 15 new sequences, which included three species: *C. qingyuanensis*, *C. peliosanthifolia* and *C. scaposa*. The remaining sequences were obtained from the GenBank public database at the National Center for Biotechnology Information (NCBI) (Table 1).

Two nuclear DNA regions (ETS 1f and ITS) and three chloroplast DNA regions (*matK*, *ndhF* and *rps16*) (at least three genes) were used for the phylogenetic analysis. The amplified primers followed Starr et al. (2003) for ETS 1f, White et al. (1990) and Blattner (1999) for ITS, Gilmour et al. (2013) for *matK* and *ndhF* and Starr et al. (2015) for *rps16*.

The sequences were aligned using the online version of MAFFT (Katoh et al. 2019). The ETS 1f, ITS, *matK*, *ndhF* and *rps16* were combined in eight Sequence Matrix (combined nDNA-cpDNA; combined nDNA; combined cpDNA; ETS 1f; ITS; *matK*; *ndhF*; and *rps16*). PhyloSuite (Zhang et al. 2020) software was used to generate the Maximum Likelihood (ML) tree and Bayesian (BI) trees, with *Eriophorum vaginatum* set as the outgroup (Starr et al. 2015). ML analysis was conducted using IQ-TREE (Nguyen et al. 2015) with 5000 standard non-parametric bootstrap replicates. BI analysis was performed using MrBayes 3.2.6 (Ronquist et al. 2012), two independent parallel chains and 5,000,000 generations

**Table 1.** Genbank numbers for samples used and in combined ITS, ETS 1f, *matK*, *ndhF*, and *rps16* analyses.

Sect.	Species	ITS	ETS 1f	<i>matK</i>	<i>ndhF</i>	<i>rps16</i>
Sect. <i>Siderostictae</i> s.s.	<i>Carex adrienii</i>	KP273628	KP273594	KP273663	KP273717	KP273771
Sect. <i>Siderostictae</i> s.s.	<i>Carex geographica</i>	KX722473	/	KX722479	KX722485	KX722491
Sect. <i>Siderostictae</i> s.l.	<i>Carex glossostigma</i>	MN762656	/	KP273686	KP273740	/
Sect. <i>Siderostictae</i> s.l.	<i>Carex grandiligulata</i>	MW459022	MW458991	MW459089	/	/
Sect. <i>Siderostictae</i> s.s.	<i>Carex kucyniakii</i>	KP273651	KP273617	KP273695	KP273749	KP273805
Sect. <i>Siderostictae</i> s.s.	<i>Carex pachygyna</i>	DQ998936	DQ998882	MW459090	/	/
Sect. <i>Siderostictae</i> s.s.	<i>Carex peliosanthifolia</i>	OR450685	OR463437	OR464515	OR464518	OR464521
Sect. <i>Siderostictae</i> s.s.	<i>Carex qingyuanensis</i>	OR450686	OR463436	OR464514	OR464517	OR464520
Sect. <i>Siderostictae</i> s.s.	<i>Carex scaposa</i>	OR450687	OR463438	OR464516	OR464519	OR464522
Sect. <i>Siderostictae</i> s.l.	<i>Carex siderosticta</i>	KP273658	KP273624	KJ513592	KJ513499	KP273817
Sect. <i>Siderostictae</i> s.s.	<i>Carex thinii</i>	KX722474	KX722468	KX722480	KX722486	KX722492
Sect. <i>Siderostictae</i> s.s.	<i>Carex tsiangii</i>	KU496610	KU377556	KU496590	/	/
Sect. <i>Siderostictae</i> s.l.	<i>Carex wuyishanensis</i>	MW459024	MW458993	MW459091	/	/
sect. <i>Hypolytroides</i>	<i>Carex hypolytroides</i>	KP273647	KP273612	KP273690	KP273744	KP273800
sect. <i>Hypolytroides</i>	<i>Carex moupinensis</i>	KP273653	KP273619	KP273699	KP273753	KP273809
Setigera Clade	<i>Carex baccans</i>	KP273632	KP273598	KP273669	KP273723	KP273778
Setigera Clade	<i>Carex myosurus</i>	KP273654	KP273620	KP273700	KP273754	KP273810
Decora Clade	<i>Carex cruciata</i>	KP273637	KP273603	KP273676	KP273730	KP273787
Decora Clade	<i>Carex filicina</i>	KP273642	KP273608	KP273682	KP273736	KP273793
Esquiroliana Clade	<i>Carex esquiroliana</i>	MN762053	MN761064	MN763585	/	/
Outgroup	<i>Eriophorum vaginatum</i>	AH012952.2	AH012952.2	KJ513615	KJ513522	KP273830



with sampling once every 100 generations. The first 25% of trees from all runs were discarded as burn-in. Finally, ITOL (Ivica and Peer 2019) software was used for tree visualisation and refinement.

## Taxonomic treatment

***Carex qingyuanensis* Y.L. Li & H.F. Chen, sp. nov.**

urn:lsid:ipni.org:names:77340693-1

清远薑草


**Type.** CHINA. Guangdong Province, Qingyuan City, Qingxin District, Bijia Mountain Forest Farm, 23°49'45"N, 113°03'07"E, 600 m elev., in the forest, on the rocks, 18 November 2022, *Li Yali & Chen Hongfeng* LYL0012 (holotype: IBSC; isotype: IBSC) (Figs 1, 2).

**Diagnosis.** The new species is similar to *Carex peliosanthifolia* F. T. Wang & Tang ex P. C. Li, but differs by having inflorescence branches racemose, single (rarely binate or ternate), one (rarely two or three) spiked, (vs. binate or ternate, one-three spiked), male part of spikes short-cylindrical or linear-cylindrical and slightly or much longer than female part (vs. just male part short-cylindrical and slightly longer than female part); style base thickened (vs. not thickened); beak short and slightly curved (vs. long and excurved).


**Description.** Perennial herbs. Rhizome stoloniferous, woody. Culms lateral, trigonous, loosely pubescent, base with brown sheaths, 15–60 cm tall. Leaves basal and cauline; basal leaves 1(2–3) tufted on rhizome node; petiole 15–25 cm long, folded, glabrous; blades narrowly elliptical, 18–30 × 2.5–5 cm, glabrous or scabrid on abaxial veins, replicate, base attenuate, apex acuminate; cauline leaves pale greenish-white with dense brown spots and short lines, spathe-like, pubescent. Involucral bracts spathe-like. Panicle compound; inflorescence branches racemose, 5–8 branched, single (rarely binate or ternate), 1 (rarely 2 or 3) spiked; peduncles of inflorescence branches tenuous, 0.5–8 cm long, densely pubescent; bractlets glume-like, ovate-oblong, ca. 3.5 mm long. Spikes bisexual, densely flowered, androgynous; spikes 6–15 mm long, male part short-cylindrical to linear-cylindrical, slightly or much longer than female part, with ca. 15–40 staminate flowers; female part with 6–25 pistillate flowers. Staminate glumes pale yellow laterally with dense spots and short lines, pale green at middle, ovate-lanceolate, ca. 3 mm long, membranous, 3-veined, apex acuminate; pistillate glumes similar to staminate glumes. Stamens 3, filaments basally connate, 0.5–3 mm long, longer or remarkably shorter than staminate glumes; anther yellow, 1 × 0.2 mm, pollen 0.2 mm wide; perigynium pale yellowish-white with brown spots and short lines, horizontally patent, elliptical, trigonous, 2.6–3.3 mm long, membranous, glabrous, with many raised veins, base subrounded, apex attenuate into a slightly curved beak, ca. 0.5 mm long. Nutlets brown at maturity, tightly enveloped, elliptical, trigonous or base obliquely truncate, 1–2 mm long; style suberect, base thickened; stigmas 3.

**Phenology.** Flowering from August to November. Fruiting from December to February.

**Etymology.** The term “qingyuanensis” originates from the location where the type specimen was collected.



**Figure 1.** **A** *Carex qingyuanensis* **B** abaxial and adaxial surface of leaf blade **C** culm **D, E** spikes **F** adaxial surface of pistillate glume **G** abaxial surface of pistillate glume **H** adaxial surface of staminate glume **I** abaxial surface of staminate glume **J, K** staminate flower **L** anther **M** perigynium **N** stigmas **O, P** nutlets **Q** the transection of nutlet. Drawn by Mrs. Liu Yunxiao based on Li Yali & Chen Hongfeng LYL0012.



**Figure 2.** **A** habitat of *Carex qingyuanensis* **B** infructescence of *C. qingyuanensis* **C** infructescence of *C. peliosanthifolia* **D** inflorescence of *C. qingyuanensis* **E** inflorescence of *C. peliosanthifolia* **F** spikes of *C. qingyuanensis* **G** spike of *C. peliosanthifolia* **H** short beak of *C. qingyuanensis* **I** long beak of *C. peliosanthifolia* **J** immature nutlets + base thickened style of *C. qingyuanensis* **K** immature nutlets + unthickened style of *C. peliosanthifolia*.


**Distribution and habitat.** *Carex qingyuanensis* is known only from Bijia Mountain Forest Farm, Qingxin District, Qingyuan City, Guangdong Province, China. It grows on rocky terrain within the forest at an elevation of 600 m (Fig. 2A).

**Conservation status.** Currently, *Carex qingyuanensis* is only known from its type locality, Bijia Mountain Forest Farm, Qingxin District, Qingyuan City, Guangdong Province, China, which covers an area of 2646.11 ha. Based on the IUCN Red List Criteria (IUCN 2022), the species could be assessed as Endangered (EN) or Vulnerable (VU). However, at present, it is more appropriate to classify it as Data Deficient (DD) due to the absence of field surveys conducted on the populations of this species.

**Additional specimens examined.** CHINA. Guangdong: Qingyuan City, Qingxin District, Bijia Mountain Forest Farm, 18 November 2022, Li Yali & Chen Hongfeng LYL0013 (IBSC); Qingyuan City, Qingxin District, Bijia Mountain Forest Farm, 1 December 2022, Li Yali & Chen Hongfeng LYL0014 (IBSC).

Results

The phylogenetic trees inferred from ML and BI shared an identical topology, while BI showed higher support values. Matrices of combined nDNA-cpDNA (Fig. 3) and combined nDNA (Suppl. material 1: fig. S1) yielded similar topologies, but slight differences in phylogenetic relationships within the sect. *Siderostictae* s.s. and the relationship between sect. *Hypolytroides* and sect. *Siderostictae* s.s.. Except for combined cpDNA (Suppl. material 1: fig. S4), *matK* (Suppl. material 1: fig. S5), and *ndhF* (Suppl. material 1: fig. S6) matrices, species within sect. *Siderostictae* s.s. consistently formed a single clade (Fig. 1, Suppl. material 1: figs S1–S3,



**Figure 3.** Phylogenetic relationships of 21 species by combined nDNA-cpDNA matrices. *Eriophorum vaginatum* was set as outgroup. The phylogenetic tree was constructed by MrBayes (BI). The bootstrap values are represented at nodes, red circles indicate bootstrap values of 90–100%, while the rest are marked with numbers.



S7). As indicated by the study conducted by Villaverde et al. (2020) and Global Carex Group (2021), analyses performed on the combined nDNA-cpDNA matrices effectively illustrate the phylogenetic relationship among 21 species (Fig. 3). *Carex baccans*, *C. myosurus*, *C. cruciata*, *C. filicina*, and *C. esquioliana* are part of the core *Carex* Clade, while species in sect. *Hypolytroides* and sect. *Siderostictae* s.s. (Global Carex Group 2021) are categorized under the *Siderosticta* clade according to Villaverde et al. (2020). The phylogenetic trees (Fig. 3) strongly reinforce the intrageneric relationships within *Siderosticta* clade and core *Carex* Clade. Furthermore, sect. *Hypolytroides* is identified as sister group to sect. *Siderostictae* s.s. (Fig. 3) remains consistent with the findings of both Villaverde et al. (2020) and Global Carex Group (2021) studies. Based on the topology, among the 13 species within sect. *Siderostictae* s.s., *C. qingyuanensis* shows closer phylogenetic relationship to *C. scaposa* rather than to *C. peliosanthifolia*.

## Discussion

According to the classification by Dai et al. (2000) and Global Carex Group (2021), *Carex qingyuanensis* belongs to sect. *Siderostictae* s. s. (traditionally placed in sect. *Hemiscaposae*) (subg. *Siderosticta*). Morphologically, it shares similarities with *C. peliosanthifolia* in having basal leaves tufted, fan-shaped; panicle compound, branches not only single and perigynium with many raised veins, but differs by having inflorescence branches racemose, single (rarely binate or ternate), one (rarely two or three) spiked, (vs. binate or ternate, 1–3-spiked), male part of spikes short-cylindrical or linear-cylindrical and slightly or much longer than the female part (vs. just male part short-cylindrical and slightly longer than female part); style base thickened (vs. not thickened) and beak short and slightly curved (vs. long and excurved) (Dai et al. 2000, 2010). Morphological comparisons of *C. qingyuanensis* and *C. peliosanthifolia* are summarised in Table 2. Furthermore, an updated identification key to Chinese species in the traditionally sect. *Hemiscaposae* is provided below.

### Key to the species of *Carex* sect. *Siderostictae* s. s. (traditional in sect. *Hemiscaposae*) in China

- 1 Ten or more spikes on each branch; leaves flat; perigynium with 2 lateral veins on adaxial surface .....2
- Fewer than 10 spikes on each branch; leaves fan-shaped; perigynium with many raised veins.....7
- 2 Panicle compound, several branched; culms rigid.....3
- Panicle simple, only one terminal branched or 1–2 lateral branched; culms flaccid.....6
- 3 Inflorescence branches paniculate, 10–20 spikes; male part of spikes oblong-cylindrical.....***C. scaposa***
- Inflorescence branches subcorymbose, a few spikes; male part of spike circular or oblong .....4
- 4 Leaves linear or linear-oblong; perigynium ovate, beak ca. 1/4 length of perigynium..... ***C. liouana***
- Leaves narrowly elliptical to elliptical-linear; perigynium elliptical, beak slightly shorter than 1/2 length of perigynium .....5



- 5 Leaves margin glabrous, scabrid abaxially; petiole glabrous; nutlets ovate ..... **C. adrienii**
- Leaves margin densely ciliate, glabrous on both surfaces or scabrid or densely hairy abaxially; petiole hairy; nutlets elliptical..... **C. densifimbriata**
- 6 Leaves narrowly elliptical to linear-elliptical, margins densely replicate; male part of spike circular to oblong, 2.5–4 mm long, ca. 3 mm wide; culms loosely hairy..... **C. lingii**
- Leaves band shape, margins flat; male part of spikes circular-cylindrical, 3–5 mm long, ca. 1 mm wide; culms loosely hairy and then glabrescent... ..... **C. ypsilandrifolia**
- 7 Branches single, rarely binate; basal leaf single..... **C. kucyniakii**
- Branches not only single; basal leaves 1–3 tufted ..... **8**
- 8 Branches binate or ternate. male part of spikes short-cylindrical ..... **C. peliosanthifolia**
- Branches single (binate or ternate). Male part of spikes short-cylindrical or linear-cylindrical..... **C. qingyuanensis**

Table 2. Morphological comparison among *Carex qingyuanensis*, *Carex peliosanthifolia*, and *Carex scaposa*.

Character	<i>C. qingyuanensis</i>	<i>C. peliosanthifolia</i>	<i>C. scaposa</i>
Inflorescence branches	Single ( rarely binate or ternate), one ( rarely two or three) spiked	Binate or ternate, one-three spiked;	Panicle compound
Spikes	Long, male part short-cylindrical or linear-cylindrical and slightly or much longer than female part	Short, male part short-cylindrical and slightly longer than female part	Male part of spike linear-lanceolate, usually shorter than female part
Pistil	Style base thickened	Style base not thickened	Style base not or slightly thickened
Perigynium	Apex attenuate into a slightly curved short beak	Apex attenuate into an excurved long beak	Apex contracted gradually into beak of medium length

Acknowledgements

We would like to express our gratitude to Hongying Zhou of Liangtian Agriculture and Forestry Technology Co., LTD, Qingyuan, Guangdong, China and Zhongcai Fan (IBSC), for their assistance in the plant management of *Carex qingyuanensis*. We appreciate the facilitation provided by the National Wild Plant Germplasm Resource Center. We also extend our appreciation to Rimei Zeng, Weifang Lu and Zhaosheng Pang for their collaboration in the field investigation of *Carex peliosanthifolia* and *Carex scaposa*. Additionally, we would like to thank Dr. Yunfei Deng (IBSC) for reviewing the article and Mrs. Yunxiao Liu (IBSC) for preparing the line drawing.

Additional information

Conflict of interest

The authors have declared that no competing interests exist.

Ethical statement

No ethical statement was reported.

Funding

This study received financial support from the Key Research and Development Program of Guangdong Province (2020B1111530004) and the Guangzhou Science and Technology Project.

## Author contributions

Hong-Feng Chen & Ya-Li Li: Conceptualization, Methodology; Hong-Feng Chen, Ya-Li Li, Jin-Chu Luo, Li-Ting Zou, Ming-Xia Li & Qiu-Gen Zeng: Field investigation, Materials collection; Ya-li Li : Data analyses and visualization; Ya-li Li: manuscript preparation; Shuang-Wen Deng & Hong-feng Chen: manuscript revision. All authors have read and approved the manuscript.

## Author ORCIDs

Ya-Li Li  <https://orcid.org/0000-0003-4667-5241>

Shuang-Wen Deng  <https://orcid.org/0000-0002-1595-2974>

Hong-Feng Chen  <https://orcid.org/0000-0002-8415-3260>

## Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

## References

- Blattner FR (1999) Direct amplification of the entire ITS region from poorly preserved plant material using recombinant PCR. *Biotechniques* 27(6): 1180–1186. <https://doi.org/10.2144/99276st04>
- Dai LK, Liang SY, Li PC, Tang YC (2000) *Carex* (Cyperaceae-Carioideae). In: Dai LK, Liang SY (Eds) *Flora Reipublicae Popularis Sinicae* 12. Science Press, Beijing, 56–528.
- Dai LK, Liang SY, Zhang SR, Tang YC, Koyama T, Tucker GC (2010) *Carex* Linnaeus. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* 23. Science Press, Beijing & Missouri Botanical Garden Press, St. Louis, 285–461.
- Deng YF (2014) *Carex longicolla* (Cyperaceae), a new sedge from China. *Phytotaxa* 178(3): 181–188. <https://doi.org/10.11646/phytotaxa.178.3.3>
- Ford BA, Starr JR, Thanh NT, Tài VA (2017) Two extraordinary new pseudopetiolate *Carex* Species from Vietnam (sect. *Hemiscaposae*, Cyperaceae). *Systematic Botany* 42(3): 402–417. <https://doi.org/10.1600/036364417X695998>
- Frodin DG (2004) History and concepts of big plant genera. *Taxon* 3(3): 753–776. <https://doi.org/10.2307/4135449>
- Global *Carex* Group (2016) Megaphylogenetic specimenlevel approaches to the *Carex* (Cyperaceae) phylogeny using ITS, ETS, and matK sequences: implications for classification. *Systematic Botany* 41(3): 500–518. <https://doi.org/10.1600/036364416X692497>
- Global *Carex* Group (2019) A tale of worldwide success: Behind the scenes of *Carex* (Cyperaceae) biogeography and diversification. *Journal of Systematics and Evolution* 57(6): 695–718. <https://doi.org/10.1111/jse.12549>
- Global *Carex* Group (2021) A framework infrageneric classification of *Carex* (Cyperaceae) and its organizing principles. *Journal of Systematics and Evolution* 59(4): 726–762. <https://doi.org/10.1111/jse.12722>
- Gilmour CN, Starr JR, Naczi RFC (2013) *Calliscirpus*, a new genus for two narrow endemics of the California Floristic Province, *C. criniger* and *C. brachythrix* sp. nov. (Cyperaceae). *Kew Bulletin* 68(1): 85–105. <https://doi.org/10.1007/s12225-012-9420-2>
- IUCN (2022) Guidelines for using the IUCN Red List Categories and Criteria. Version 15.1. Prepared by the Standards and Petitions Subcommittee. IUCN Red List of Threatened Species. [accessed on July 2022]

- Ivica L, Peer B (2019) Interactive Tree Of Life (iTOL) v4: Recent updates and new developments. *Nucleic Acids Research* 47(W1): W256–259. <https://doi.org/10.1093/nar/gkz239>
- Jin XF (2009) *Carex yunyiana* sp. nov. (Cyperaceae) from zhejiang, China. *Nordic Journal of Botany* 27(4): 344–347. <https://doi.org/10.1111/j.1756-1051.2009.00452.x>
- Jin XF (2017) Taxonomic revision of *Carex chungii* (Cyperaceae; sect. *Mitratae*) and allied species. *Phytotaxa* 317 (1): 029–041. <https://doi.org/10.11646/phytotaxa.317.1.3>
- Jin XF, Chen WJ (2015) *Carex huangshanica* (sect. *Rhomboidales*, Cyperaceae), a new species from anhui, eastern China. *Phytotaxa* 233(3): 281–287. <https://doi.org/10.11646/phytotaxa.233.3.5>
- Jin XF, Zheng CZ (2010) *Carex yandangshanica* sp. nov. (Cyperaceae; C. sect. *Rhomboidales*) from Zhejiang, China. *Nordic Journal of Botany* 28: 709–712. <https://doi.org/10.1111/j.1756-1051.2010.00817.x>
- Jin XF, Zheng CZ, Ding BY (2004) New taxa of *Carex* (Cyperaceae) from Zhejiang, China. *Acta Phytotaxonomica Sinica* 42(6): 541–550.
- Jin XF, Zhao YJ, Zheng CZ, Zhang HW (2011) *Carex zhejiangensis* sp. nov. (Cyperaceae) from Zhejiang, eastern China. *Nordic Journal of Botany* 29(1): 63–70. <https://doi.org/10.1111/j.1756-1051.2010.00977.x>
- Jin SH, Zhao YJ, Shen TT, Wu DF, Jin XF (2012a) *Carex tenuirostrata* (Cyperaceae), a new species of section *Confertiflorae* from Zhejiang, China. *Brittonia* 64(3): 325–329. <https://doi.org/10.1007/s12228-012-9236-2>
- Jin XF, Simpson DA, Zheng CZ, Sun L, Zhang HW (2012b) *Carex paracheniana* (*Carex* sect. *Rhomboidales*, Cyperaceae), a new species from Guangxi and Guizhou, China. *Systematic Botany* 37(4): 929–937. <https://doi.org/10.1600/036364412X656617>
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics* 20(4): 1160–1166. <https://doi.org/10.1093/bib/bbx108>
- Li YY, Chen YP, Jiang LQ, Liu ED, Luo Y, Peng H (2022) *Carex malipoensis* (Cyperaceae), a new species from southeast Yunnan, China. *PhytoKeys* 188: 19–29. <https://doi.org/10.3897/phytokeys.188.75401>
- Lu YF, Jin XF (2018) Notes on *Carex* (Cyperaceae) from China (V): new species, combination and subspecies from Zhejiang. *Phytotaxa* 372(3): 203–211. <https://doi.org/10.11646/phytotaxa.372.3.3>
- Lu YF, Jin XF (2022) Notes on *Carex* (Cyperaceae) from China (VIII): Five new species and a new variety from southern and south-western China. *PhytoKeys* 188: 31–47. <https://doi.org/10.3897/phytokeys.188.77776>
- Lu ZC, Su YL, Lu YF, Jin XF (2020) *Carex pingleensis* (*Carex* sect. *Mitratae*), a new species of Cyperaceae from Guangxi, China. *Taiwania* 65(3): 391–395. <https://doi.org/10.6165/tai.2020.65.391>
- Lu YF, Jin XF, Yu MJ (2022) *Carex borealifujianica* (Cyperaceae), a new species of the core *Carex* clade from Fujian, southeastern China. *PLoS ONE* 17(3): e0264419. <https://doi.org/10.1371/journal.pone.0264419>
- Lunkai D, Liang S, Liang SJ, Zhang S, Tang Y, Tang YC, Koyama T, Tucker GC (2010) *Carex*. In: Wu ZY, Raven PH, Hong DY (Eds) *Flora of China* vol. 23. Science Press and Missouri Botanical Garden Press, Beijing, 285–461.
- Nelmes E (1955) The genus *Carex* in Indo-China including Thailand and Lower Burma. *Mémoires du Muséum national d'histoire naturelle. Série B. Botanique* 4: 83–182. <https://www.biodiversitylibrary.org/part/283006>



- Nguyen LT, Schmidt HA, Haeseler A, Minh BQ (2015) IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* 32(1): 268–274. <https://doi.org/10.1093/molbev/msu300>
- Raymond M (1959) Carices Indochinenses necnon Siamenses. *Mémoires de Jardin Botanique de Montréal* 53: 1–125.
- Ronquist FM, Teslenko P, van der Mark DL, Ayres A, Darling S, Höhna B, Liu L, Suchard MA, Huelsenbeck JP (2012) MRBAYES 3.2: Efficient Bayesian phylogenetic inference and model selection across a large model space. *Systematic Biology* 61(3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Song C, Wang H, Jin XF (2008) A new subspecies of *Carex arisanensis* (Cyperaceae) from zhejiang, China. *Annales Botanici Fennici* 45(2): 156–158. <https://doi.org/10.5735/085.045.0212>
- Starr JR, Harris SA, Simpson DA (2003) Potential of the 5' and 3' Ends of the Intergenic Spacer (IGS) of rDNA in the Cyperaceae: New Sequences for Lower-Level Phylogenies in Sedges with an Example from *Uncinia* Pers. *International Journal of Plant Sciences* 164(2): 213–227. <https://doi.org/10.1086/346168>
- Starr JR, Janzen FH, Ford BA (2015) Three new, early diverging *Carex* (Cariceae, Cyperaceae) lineages from East and Southeast Asia with important evolutionary and biogeographic implications. *Molecular Phylogenetic Evolution* 88: 105–120. <https://doi.org/10.1016/j.ympev.2015.04.001>
- Su SW (2009a) Additional notes on the genus *Carex* Linn. from Anhui, China. *Journal of Anhui Agricultural University* 36: 557–563. <https://doi.org/10.13610/j.cnki.1672-352x.2009.04.001>
- Su SW (2009b) *Carex kobresiformis*, a particular species of *Carex* from China. *Journal of Anhui University* 33(6): 83–86.
- Uzma Jiménez-Mejías P, Amir R, Hayat MQ, Hipp AL (2019) Timing and ecological priority shaped the diversification of sedges in the Himalayas. *PeerJ* 7: e6792. <https://doi.org/10.7717/peerj.6792>
- Villaverde T, Jiménez-Mejías P, Luceño M, Waterway MJ, Kim S, Lee B, Rincón-Barrado M, Hahn M, Maguilla E, Roalson EH, Hipp AL, Global Carex Group (2020) A new classification of *Carex* subgenera supported by a HybSeq backbone phylogeny. *Botanical Journal of the Linnean Society* 194(2): 141–163. <https://doi.org/10.1093/botlinnean/boaa042>
- Wang QL, Yang HB, Deng YF, Wang ZN, Liu GD (2012) *Carex longipetiolata* (Cyperaceae), a new sedge from Hainan, China. *Phytotaxa* 75(1): 65–69. <https://doi.org/10.11646/phytotaxa.75.1.6>
- Waterway MJ, Hoshino T, Masaki T (2009) Phylogeny, species richness, and ecological specialization in Cyperaceae tribe Cariceae. *Botanical Review* 75(1): 138–159. <https://doi.org/10.1007/s12229-008-9024-6>
- Weng DM, Zhang HW, Xu SF, Jin XF (2009) *Carex qingliangensis* sp. nov. (Cyperaceae) from zhejiang, eastern china. *Nordic Journal of Botany* 27(1): 7–10. <https://doi.org/10.1111/j.1756-1051.2009.00394.x>
- White TJ, Bruns T, Lee S, Taylor WJ, Innis MA, Gelfand DH, Sninsky J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and amplifications*, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>

- Yang SZ, Chen CF (2005) *Carex scaposa* C. B. Clarke (Cyperaceae): A new record to the flora of Taiwan. *Taiwania* 50(3): 227–233. [https://doi.org/10.1016/S0011-2240\(05\)00046-5](https://doi.org/10.1016/S0011-2240(05)00046-5)
- Yang HB, Liu GD (2020) *Carex ledongensis* (Cyperaceae), a new sedge from Hainan Isl. of South China. *Phytotaxa* 461(2): 72–78. <https://doi.org/10.11646/phytotaxa.461.2.1>
- Yang HB, Wang QL, Bai CJ, Li XX, Liu GD (2014) *Carex diaoluoshanica* (*Carex* sect. *Lageniformes*, Cyperaceae), a New Species from Hainan, China. *PLoS ONE* 9(6): e97658. <https://doi.org/10.1371/journal.pone.0097658>
- Yang HB, Li XX, Chang JB, Wang WQ, Wang QL, Yu DG, Dong RS, Liu GD (2015a) *Carex procumbens* (*Carex* sect. *Rhomboidales*, Cyperaceae), a new species from Hainan, China. *Phytotaxa* 201(3): 207–213. <https://doi.org/10.11646/phytotaxa.201.3.4>
- Yang HB, Li XX, Wang WQ, Bai CJ, Liu GD (2015b) *Carex jianfengensis* (*Carex* sect. *Rhomboidales*, Cyperaceae), a new species from Hainan, China. *PLoS ONE* 10(9): e0136373. <https://doi.org/10.1371/journal.pone.0136373>
- Yang HB, Li XX, Bai CJ, Wang WQ, Liu GD (2016) *Carex concava* (*Carex* sect. *Rhomboidales*, Cyperaceae), a new species from Hainan, China. *Phytotaxa* 283(2): 155–162. <https://doi.org/10.11646/phytotaxa.283.2.4>
- Yang HJ, Xu H, Deng YF, Li YD (2017) *Carex xueyingiana* (Cyperaceae), a new species from Hainan Island, China. *Phytotaxa* 308(2): 239–248. <https://doi.org/10.11646/phytotaxa.308.2.5>
- Yu LY, Huang YF, Xu WB, Deng YF (2012) *Carex trongii* K. K. Nguyen, a newly recorded species of Cyperaceae from China. *Journal of Tropical and Subtropical Botany* 20(4): 403–406. <https://doi.org/10.3969/j.issn.1005-3395.2012.04.014>
- Zhang SR, Zhang J, Liu ZY, Qu S, Han RG (2018) *Carex nodosa* (Cyperaceae), a new species of *C.* sect. *Occlusae* from southwest China. *Nordic Journal of Botany* 36(8): 1–5. <https://doi.org/10.1111/njb.01900>
- Zhang D, Gao F, Jakovlić I, Zou H, Zhang J, Li WX, Wang G (2020) PhyloSuite: An integrated and scalable desktop platform for streamlined molecular sequence data management and evolutionary phylogenetics studies. *Molecular Ecology Resources* 20(1): 348–355. <https://doi.org/10.1111/1755-0998.13096>
- Zhang XX, Yao ZM, Meng HL, Deng YF (2021) *Carex maolanensis* (Cyperaceae), a new species from Guizhou, China. *Phytotaxa* 525(2): 137–146. <https://doi.org/10.11646/phytotaxa.525.2.4>
- Zhao YJ, Shen TT, Zheng Y, Chen ZL, Jin XF (2011) *Carex dapanshanica* sp. nov. (*C.* sect. *Mitratae*, Cyperaceae) from Zhejiang, China. *Nordic Journal of Botany* 29(6): 670–673. <https://doi.org/10.1111/j.1756-1051.2011.01132.x>

## Supplementary material 1

### Phylogenetic relationships

Authors: Ya-Li Li, Shuang-Wen Deng, Jin-Chu Luo, Ming-Xia Li, Li-Ting Zou, Qiu-Gen Zeng, Hong-Feng Chen

Data type: pdf

Explanation note: **fig. S1.** Phylogenetic relationships of 21 species by combined nDNA matrices. *Eriphorum vaginatum* was set as the outgroup. The phylogenetic tree was constructed by MrBayes (BI). The bootstrap values are represented at nodes, red circles indicate bootstrap values of 90–100, while the rest are marked with numbers. The species marked in yellow belong to the Decora Clade, the ones in pink belong to the Esquiroliana Clade, the ones in green belong to the Setigera Clade, the ones in purple belong to sect. *Hypolytroides*, and those in blue belong to sect. *Siderostictae* s.s.. The ones highlighted in red belong to sect. *Siderostictae* s.s. and share similar morphological characteristics. **fig. S2.** Phylogenetic relationships of 19 species by ETS matrices. The notes are same as fig. S1. **fig. S3.** Phylogenetic relationships of 21 species by ITS matrices. The notes are same as fig. S1. **fig. S4.** Phylogenetic relationships of 21 species by combined cpDNA matrices. The notes are same as fig. S1. **fig. S5.** Phylogenetic relationships of 21 species by *matK* matrices. The notes are the same as fig. S1. **fig. S6.** Phylogenetic relationships of 16 species by *ndhF* matrices. The notes are the same as fig. S1. **fig. S7.** Phylogenetic relationships of 15 species by *rps16* matrices. The notes are the same as fig. S1.

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.241.117734.suppl1>